

COHOMOLOGY THEORIES FOR HIGHLY STRUCTURED RING SPECTRA

A. LAZAREV

ABSTRACT. This is a survey paper on cohomology theories for A_∞ and E_∞ ring spectra. Different constructions and main properties of topological André-Quillen cohomology and of topological derivations are described. We give sample calculations of these cohomology theories and outline applications to the existence of A_∞ and E_∞ structures on various spectra. We also explain the relationship between topological derivations, spaces of multiplicative maps and moduli spaces of multiplicative structures.

1. INTRODUCTION

In recent years algebraic topology witnessed renewed interest to highly structured ring spectra first introduced in [22]. To a large extent it was caused by the discovery of a strictly associative and symmetric smash product in the category of spectra in [7], [14]. This allowed one to replace the former highly technical notions of A_∞ and E_∞ ring spectra by equivalent but conceptually much simpler notions of S -algebras and commutative S -algebras respectively.

An S -algebra is just a monoid in the category of spectra with strictly symmetric and associative smash product (hereafter referred to as the category of S -modules). Likewise, a *commutative S -algebra* is a commutative monoid in the category of S -modules.

The most important formal property of categories of S -algebras and commutative S -algebras is that both are *topological model categories* in the sense of Quillen, [26] as elaborated in [7]. That means that together with the usual structure of the closed model category (fibrations, cofibrations and weak equivalences subject to a set of axioms) these categories are topologically enriched, that is their *Hom* sets are topological spaces and the composition of morphisms is continuous. Moreover, there exist *tensors* and *cotensors* of objects with topological spaces that satisfy the usual adjunction isomorphisms which hold for cartesian products and mapping spaces in the category of topological spaces. In addition, this enrichment is supposed to be compatible with the closed model structure in the sense that an appropriate analogue of Quillen's corner axiom SM7 is satisfied.

This rich structure allows one to translate a lot of the notions of conventional homotopy theory (homotopy relation, cellular approximation, the formation of homotopy limits and colimits) in the abstract setting. The language of closed model categories will be freely used here and we refer the reader to the monograph [13] where the necessary details may be found.

The categories of S -algebras and commutative S -algebras are in some sense analogous to the unstable category of topological spaces. In these categories there are certain natural homotopy invariant theories which play the role of singular cohomology for topological spaces. In particular they are natural homes for various obstruction groups. The corresponding theory is called *topological André-Quillen cohomology* in the commutative case and *topological derivations* in the associative case. We will use the abbreviation TAQ for the former and Der for the latter. The purpose of the present paper is to give an overview of recent results on these cohomology theories. The technical level will be kept to a minimum, however we will try to outline proofs of the results we formulate, especially when these proofs are easy and conceptual. Occasionally the theorems we obtain are 'new', that is, have not appeared in print before; but in each case they are either direct extensions of the known results or could be obtained following similar patterns. Still, we tried to

put these results in different perspectives than those in the published sources; in particular the associative and commutative cases are treated as uniformly as possible.

The paper does not pretend to be comprehensive by any means. One notable omission is the work of Robinson and Whitehouse on Γ -homology of commutative rings [29] and the subsequent work of Robinson [31] where Γ -homology was used to show the existence of a commutative S -algebra structure on E_n , the so-called Morava E -theory spectrum. One should also mention the important papers [28] and [25] containing further results on Γ -homology. Our limited expertise as well as the lack of space prevented us from including these results in the survey. Another theme only briefly mentioned here is the existence of the action of the Morava stabilizer group on the spectrum E_n (the Hopkins-Miller theorem). For this and related topics we refer the reader to [27], [17] and [10].

The paper is organized as follows. In sections 2 and 3 we outline the construction and basic properties of TAQ and Der. In section 4 we investigate the problem of lifting a map of S -algebras or commutative S -algebras and introduce the notion of a *topological singular extension*. In section 5 we discuss an alternative construction of TAQ and Der as *stabilizations* of appropriate forgetful functors. Section 6 presents calculations of TAQ and Der of the Eilenberg-MacLane spectrum $H\mathbb{F}_p$. In section 7 we show how the developed technology could be used to produce MU -algebra structures on many complex oriented spectra. Here MU is the spectrum of complex cobordisms. In section 8 we discuss the relationship between spaces of algebra maps and Der. In section 9 we make a link between Der and spaces of multiplicative structures on a given spectrum.

Notation and conventions. The paper is written in the language of S -modules of [7] and we adopt the terminology of the cited reference. There is one exception: we use the terms ‘cofibrant’ and ‘cofibration’ for what [7] calls ‘ q -cofibrant’ and ‘ q -cofibration’. This is because we never have the chance to use cofibrations in the classical sense (maps that satisfy the homotopy extension property). Except for Section 6 we work over an arbitrary (but fixed) cofibrant commutative S -algebra R , smash products $\wedge$ and function objects $F(-, -)$ mean $\wedge_R$ and $F_R(-, -)$; when we use a different ‘ground’ S -algebra A this is indicated by a subscript such as $\wedge_A$ and $F_A(-, -)$. The topological space of maps in a topological category $\mathcal{C}$ is denoted by $\text{Map}_{\mathcal{C}}$. The category of unbased topological spaces is denoted by Top and that of based spaces - by Top_* . For an S -algebra A the category of (left) A -modules is denoted by $\mathcal{M}_A$. The category of A -bimodules (which is the same as the category of $A \wedge A^{op}$ -modules where A^{op} is the R -algebra A with opposite multiplication) will be denoted by $\mathcal{M}_{A-A}$. The category of R -algebras is denoted by $\mathcal{C}_R^{ass}$ and that of *commutative* R -algebras - by $\mathcal{C}_R^{comm}$. The homotopy category of a closed model category $\mathcal{C}$ is denoted by $h\mathcal{C}$ so, for example $h\mathcal{C}_R^{comm}$ is the homotopy category of commutative R -algebras. For an R -algebra A we will denote by $[-, -]_{A-mod}$ and $[-, -]_{A-bimod}$ the sets of homotopy classes of A -module or A -bimodule maps respectively. The field consisting of p elements is denoted by $\mathbb{F}_p$.

Acknowledgement. I would like to express my sincere gratitude to Andy Baker and Birgit Richter for doing such a wonderful job of organizing the Workshop on Structured Ring Spectra in Glasgow in January 2002. I am also thankful to Maria Basterra for explaining to me her joint work with Mike Mandell on TAQ and to Stefan Schwede for pointing out that Theorem 9.2 is a direct consequence of Theorem 8.1. Thanks are also due to Mike Mandell, Paul Goerss, Nick Kuhn and Bill Dwyer for useful discussions and comments made at various times.

2. TOPOLOGICAL ANDRÉ-QUILLEN COHOMOLOGY FOR COMMUTATIVE S -ALGEBRAS

Let A be a commutative R -algebra which without loss of generality will be assumed to be cofibrant. Denote by $\mathcal{C}_{R/A}^{comm}$ the category of commutative R -algebras over A . An object of $\mathcal{C}_{R/A}^{comm}$ is a commutative R -algebra B supplied with an R -algebra map $B \rightarrow R$ (an augmentation). A morphism

between two such objects B and C is the following commutative diagram in $\mathcal{C}_R^{comm}$:

$$\begin{array}{ccc} B & \xrightarrow{\quad} & C \\ & \searrow & \swarrow \\ & A & \end{array}$$

Then $\mathcal{C}_{R/A}^{comm}$ inherits a topological model category structure from $\mathcal{C}_R^{comm}$ so that a map in $\mathcal{C}_{R/A}^{comm}$ is a cofibration if it is so considered as a map in $\mathcal{C}_R^{comm}$. Note that in the case $R = A$ the category $\mathcal{C}_{R/A}^{comm} = \mathcal{C}_{A/A}^{comm}$ is pointed and therefore is enriched over the category of pointed topological spaces.

Let us denote by $\mathcal{N}_A$ the category of *nonunital* commutative A -algebras. An object in this category is an A -module M together with a strictly associative multiplication map $M \wedge_A M \rightarrow M$. The morphisms in $\mathcal{N}_A$ are defined in the obvious fashion. Following [1] we will refer to an object of $\mathcal{N}_A$ as an A -NUCA. It is not hard to prove that $\mathcal{N}_A$ has a topological model structure where weak equivalences are those maps which are weak equivalences on underlying A -modules. The fibrations then are the maps that are fibrations of underlying A -modules and the cofibrations are the maps which have the LLP with respect to the acyclic fibration. Then we could form the homotopy category $h\mathcal{N}_A$ of $\mathcal{N}_A$.

We want to show that the categories $h\mathcal{C}_{A/A}^{comm}$ and $h\mathcal{N}_A$ are equivalent. In fact, more is true, see Proposition 2.1.

Let $K : \mathcal{N}_A \rightarrow \mathcal{C}_{A/A}^{comm}$ denote the functor which assigns to an A -NUCA M the commutative A -algebra $A \vee M$ with multiplication

$$(A \vee M) \wedge_A (A \vee M) \cong A \vee M \vee M \vee M \vee M \wedge_A M \rightarrow A \vee M$$

given by the obvious maps on the first three wedge summands and by the multiplication of M on the last one.

Clearly the A -algebra $A \vee M$ can be considered as an object in $\mathcal{C}_{A/A}^{comm}$ via the canonical projection map $A \vee M \rightarrow A$. In other words the functor K lands in fact in the category $\mathcal{C}_{A/A}^{comm}$.

Now let B be an A -algebra over A and denote by I its ‘augmentation ideal’, i.e. the fibre of the augmentation map $B \rightarrow A$. Then $I(B)$ is naturally an A -NUCA and we can consider I as a functor from $\mathcal{C}_{A/A}^{comm}$ to $\mathcal{N}_A$.

Then we have the following

Proposition 2.1. *The functor K is left adjoint to I . Moreover the functors K and I establish a Quillen equivalence between the categories $\mathcal{N}_A$ and $\mathcal{C}_{A/A}^{comm}$.*

To see that the functors K and I are adjoint notice that the category $\mathcal{N}_A$ is in fact a category of algebras over a certain monad $\mathbb{A}$ in A -modules. This monad is specified by $\mathbb{A}M := \bigvee_{i>0} M^{\wedge_A i} / \Sigma_i$ where Σ_i is the symmetric group on i symbols. Further note that $K(\mathbb{A}M)$ is a free commutative A -algebra on M . Therefore if $M = \mathbb{A}M$ for an A -module M we have

$$\mathcal{C}_{A/A}^{comm}(K(\mathbb{A}M), B) \cong \mathcal{M}_A(M, I(B)) \cong \mathcal{N}_A(\mathbb{A}M, B).$$

To get the adjointness isomorphism for a general A -NUCA X it suffices to notice that there is a canonical split coequalizer (the beginning of the monadic bar-construction for M):

$$\mathbb{A}\mathbb{A}X \rightrightarrows \mathbb{A}X \rightarrow M.$$

Explicitly, if $g : M \rightarrow I(B)$ is a morphism in $\mathcal{N}_A$ then its adjoint $\tilde{g} : A \vee M \rightarrow B$ is the composite

$$\tilde{g} : A \vee M \xrightarrow{id \vee g} A \vee I(B) \xrightarrow{1 \vee i} B$$

where i is the canonical map $I(B) \rightarrow B$.

Further straightforward arguments show that the functor K preserves cofibrations and acyclic cofibrations and that the adjunction described above determines an equivalence on the level of homotopy categories.

We will denote the homotopy invariant extension (also called a *total derived functor*) of the functor I by $\mathbf{R}I$. For an object B in $\mathcal{C}_{A/A}^{comm}$ denote by $\tilde{B}$ its fibrant replacement. Then $\mathbf{R}I(B) = I(\tilde{B})$. Next we define the functor of ‘taking the indecomposables’ that assigns to an A -NUCA N the A -module $Q(N)$ given by the cofibre sequence:

$$N \wedge_A N \xrightarrow{m} N \longrightarrow Q(N).$$

Here m stands for the multiplication map. The functor Q has a right adjoint functor $Z : \mathcal{M}_A \longrightarrow \mathcal{N}_A$ which is given by considering A -modules as objects in $\mathcal{N}_A$ with zero multiplication. It is easy to see that this adjunction passes to the homotopy categories. Let $\mathbf{L}Q$ denote the total derived functor of Q . Explicitly, $\mathbf{L}Q(N) = Q(\tilde{N})$ where $\tilde{N}$ is the cofibrant replacement of the A -NUCA N .

It is clear that $Q(\mathbb{A}_A) \cong N$. However for a general cofibrant A -NUCA the functor $Q(N)$ is very hard to compute. One can approach its computation as follows. The functor $\mathbb{A} : \mathcal{M}_A \longrightarrow \mathcal{M}_A[\mathbb{A}]$ from A -modules to $\mathbb{A}$ -algebras is left adjoint to the forgetful functor $U : \mathcal{M}_A[\mathbb{A}] \longrightarrow \mathcal{M}_A$. There results a monad in $\mathcal{M}_A$. Given an A -NUCA N denote by $B_*(N) = B_*(\mathbb{A}, U\mathbb{A}, UN)$ its monadic bar-construction. Explicitly B_*N is the cosimplicial A -NUCA with $B_n(N) = \mathbb{A}^{n+1}N$ the faces and codegeneracies are the standard ones, cf. [23]. The geometric realization $|B_*(N)|$ of $B_*(N)$ is weakly equivalent to N . Then one has the following

Proposition 2.2. *For a cofibrant A -NUCA N one has the following weak equivalence of A -modules: $Q(|B_*(N)|) \simeq Q(N)$.*

Despite the innocent appearance of this proposition it is not at all obvious. The problem is that the functor Q only preserves weak equivalences between cofibrant objects and $|B_*(N)|$ is not a cofibrant A -NUCA. However in the end it turns out that it is close enough to being cofibrant to give the result; much of the work [1] is devoted to overcoming this point.

Further associated to the simplicial A -NUCA $B_*(N)$ is the spectral sequence which computes $\pi_*|B_*(N)| = \pi_*N$:

$$(2.1) \quad E_{i,j}^1 = \pi_i Q(\mathbb{A}^{j+1}N) = \pi_i \mathbb{A}^j N \implies \pi_{i-j} Q(N).$$

We now have all the ingredients to define the *abelianization* functor:

$$\Omega_A^{comm} : h\mathcal{C}_{R/A}^{comm} \longrightarrow h\mathcal{M}_A.$$

Definition 2.3. Let C be a commutative R -algebra over A and M be a A -module. Then $\Omega_A^{comm}(C) := \mathbf{L}Q \circ \mathbf{R}I(C \wedge_R^{\mathbf{L}} A)$.

(Note that $C \wedge_R^{\mathbf{L}} A$ is an object of $\mathcal{C}_{A/A}^{comm}$ in an obvious way.) Of course we only defined the abelianization functor on objects, but its extension to morphisms is immediate. The main property of the abelianization functor is that it is left adjoint to the ‘square-zero’ extension functor: if M is a A -module then the square zero extension of A by M is the A -algebra over A whose underlying R -module is $A \vee M$ with the obvious multiplication. More precisely we have the following

Theorem 2.4. *There is the following natural isomorphism:*

$$h\mathcal{C}_{R/A}^{comm}(C, A \vee M) \cong h\mathcal{M}_A(\Omega_A^{comm}(C), M)$$

where C is a commutative R -algebra over A and M is a A -module.

The proof of the theorem is rather formal and relies on the already established properties of the functors I, K, Z and Q .

Remark 2.5. We would like to emphasize here that the isomorphism of Theorem 2.4 holds on the level of *homotopy* categories and is not a reflection of an adjunction between strict categories. Indeed, even though the abelianization functor could be considered as a point-set level functor its definition involves composition of right adjoint and left adjoint functors. Therefore one cannot expect any good formal properties of Ω_A^{comm} on the point-set level. The situation improves upon passing to homotopy categories because one of these functors, namely RI , becomes an equivalence.

We will sometimes need the enriched version of Theorem 2.4. The category $\mathcal{C}_{R/A}^{comm}$ is enriched over unbased topological spaces. For two objects B, A of a topological category $\mathcal{C}$ the space of maps $\text{Map}_{\mathcal{C}/A}(B, C)$ from B to C *over* A is defined from the pullback diagram

$$\begin{array}{ccc} \text{Map}_{\mathcal{C}/A}(B, C) & \longrightarrow & pt \\ \downarrow & & \downarrow \\ \text{Map}_{\mathcal{C}}(B, C) & \longrightarrow & \text{Map}_{\mathcal{C}}(B, A) \end{array}$$

where the right downward arrow is just picking the structure map $B \rightarrow A$ in $\text{Map}_{\mathcal{C}}(B, A)$. It is easy to see that the cotensor $(A \vee M)^X$ of X and $A \vee M$ in $\mathcal{C}_{R/A}^{comm}$ is weakly equivalent to $A \vee M^X$. Then for any CW -complex X we have the following isomorphisms:

$$\begin{aligned} h\mathcal{T}op(X, \text{Map}_{\mathcal{M}_C}(\Omega_A^{comm}(C), M)) &\cong h\mathcal{M}_C(\Omega_A^{comm}(C), M^X) \\ &\cong h\mathcal{C}_{R/A}^{comm}(C, A \vee M^X) \\ &\cong h\mathcal{T}op(X, \text{Map}_{\mathcal{C}_{R/A}}(C, A \vee M)). \end{aligned}$$

Therefore the topological spaces $\text{Map}_{\mathcal{C}_{R/A}}(C, A \vee M)$ and $\text{Map}_{\mathcal{M}_C}(\Omega_A^{comm}(C), M)$ are weakly equivalent.

We will denote the A -module $\Omega_A^{comm}(A) = \mathbf{L}Q \circ \mathbf{R}I(A \wedge_R^{\mathbf{L}} A)$ simply by Ω_A^{comm} so that

$$h\mathcal{C}_{R/A}^{comm}(A, A \vee M) \cong h\mathcal{M}_A(\Omega_A^{comm}, M).$$

The set $h\mathcal{C}_{R/A}^{comm}(A, A \vee M)$ could be interpreted as derivations of the commutative S -algebra A with coefficients in the A -module M . In particular we see, that the set of such derivations is an abelian group (note that apriori the set of homotopy classes of (commutative) S -algebra maps does not carry a structure of a group, let alone an abelian group).

We can now define the *topological André-Quillen cohomology* spectrum of A relative to R with coefficients in M by

$$\mathbf{TAQ}_R(A, M) := F_A(\Omega_A^{comm}, M)$$

and similarly the *topological André-Quillen homology* spectrum of A relative to R with coefficients in M :

$$\mathbf{TAQ}^R(A, M) := \Omega_A^{comm} \wedge_A M.$$

We will refer to the homotopy groups of these spectra as topological André-Quillen (co)homology of A relative to R :

$$\begin{aligned} \text{TAQ}_R^*(A, M) &:= \pi_{-*} F_A(\Omega_A^{comm}, M); \\ \text{TAQ}_*^R(A, M) &:= \pi_* \Omega_A^{comm} \wedge_A M. \end{aligned}$$

Remark 2.6. The enriched version of Theorem 2.4 gives us the weak equivalence of spaces $\text{Map}_{\mathcal{C}_{R/A}^{comm}}(B, A \vee M) \simeq \text{Map}_{\mathcal{M}_A}(\Omega_A^{comm}(C), M)$. The latter mapping space is in turn weakly equivalent to the zeroth space of the spectrum $F_A(\Omega_A^{comm}, M) = \mathbf{TAQ}_R(A, M)$. We obtain the weak equivalence

$$\text{Map}_{\mathcal{C}_{R/A}^{comm}}(B, A \vee M) \simeq \Omega^\infty \mathbf{TAQ}_R(A, M).$$

We are most interested in topological André -Quillen *cohomology* since they are related to the obstruction theory. Repeat, that for a commutative R -algebra A and an A -module M the abelian group $TAQ_R^0(A, M)$ is identified with the set of *commutative derivations* of A with values in M that is the homotopy classes of maps in $\mathcal{C}_{R/A}^{comm}$ from A into $A \vee M$. We say ‘commutative derivations’ to distinguish them from just ‘derivations’ or ‘topological derivations’ which will be considered in the next section and refer to the *bimodule* derivations in the context of associative R -algebras.

Let us now discuss the functoriality of TAQ . Since by definition $\mathbf{TAQ}_R(A, M) = F_A(\Omega_A^{comm}, M)$ we see that it is covariant with respect to the variable M . On the level of commutative derivations it can be seen as follows. A map of A -bimodules $f : M \rightarrow N$ determines a map in $\mathcal{C}_{R/A}^{comm} : A \vee M \rightarrow A \vee N$. Then a topological derivation $d : A \rightarrow A \vee M$ determines a topological derivation $f_*d : A \rightarrow A \vee M \rightarrow A \vee N$.

If $g : X \rightarrow A$ is a map of R -algebras then the composite map

$$X \xrightarrow{g} A \xrightarrow{d} A \vee M$$

is a map in $\mathcal{C}_{A/R}^{comm}$. Therefore by Theorem 2.4 it corresponds to a map of in $h\mathcal{M}_A : \Omega_A^{comm}(X) \rightarrow M$ which is the same as a map in $h\mathcal{M}_X : \Omega_X^{comm} \rightarrow M$. The latter corresponds, again by Theorem 2.4 to a commutative derivation of X with values in the X -module M . We will denote this derivation by $g^*d : X \rightarrow X \vee M$.

We can relate the topological André-Quillen cohomology to ordinary R -module cohomology as follows. For a cofibrant R -algebra A consider the composite map of R -modules

$$l_R : A \rightarrow A \vee \Omega_A^{comm} \rightarrow \Omega_A^{comm}.$$

Here the first map is the R -algebra map adjoint to the identity on Ω_A^{comm} and the second map is the projection onto the wedge summand. The map l_R induces a forgetful map from TAQ to R -module cohomology:

$$l_R^* : TAQ_R^*(A, M) = [\Omega_A^{comm}, M]_{A-mod}^* \rightarrow [\Omega_A^{comm}, M]_{R-mod}^* \rightarrow [A, M]_{R-mod}^*.$$

To describe the image of l_R^* let us introduce the notion of a *primitive operation* $E \rightarrow M$ for a R -ring spectrum E and an E -bimodule spectrum M . Namely, the map $p \in [E, M]_{R-mod}^*$ is primitive if the following diagram is commutative in $h\mathcal{M}_R$:

$$\begin{array}{ccc} E \wedge E & \xrightarrow{m} & E \\ 1 \wedge p \vee p \wedge 1 \downarrow & & \downarrow p \\ E \wedge M \vee M \wedge E & \xrightarrow{m_l \vee m_r} & M \end{array}$$

where m_l and m_r denote the left and right actions of E on M . (In other words, p is a derivation up to homotopy but we refrain from using this term to avoid confusion.) Then it is easy to see that the image of l_R^* is contained in the subspace of primitive operations in $[A, M]_{R-mod}^*$. Of course the left and right A -module spectrum structures on M coincide in the commutative case.

3. TOPOLOGICAL DERIVATIONS

In this section we construct the analogue of TAQ for not necessarily commutative S -algebras. Traditionally the André-Quillen cohomology was considered for commutative algebras only. Therefore we will reserve this term for the commutative case and call its analogue for associative S -algebras *topological derivations*. The main result we are going to describe here is the analogue of Theorem 2.4 in the context of noncommutative S -algebras. Note, however, that the construction and the proof differ considerably from the commutative case.

Let A be a cofibrant R -algebra. We define the module of differentials Ω_A^{ass} from the following homotopy fibre sequence

$$(3.1) \quad \Omega_A^{ass} \longrightarrow A \wedge A \longrightarrow A.$$

Here the second arrow is the multiplication map. Clearly Ω_A^{ass} is an A -bimodule. Note that the sequence (3.1) splits in the homotopy category of left A -modules by the map

$$A \xrightarrow{\cong} A \wedge R \xrightarrow{id \wedge 1} A \wedge A$$

and therefore the A -bimodule Ω_A^{ass} is equivalent as a left A -module to $A \wedge A/R$. Similarly Ω_A^{ass} is equivalent as a *right* A -module to $A/R \wedge A$.

Denote the category of objects over A inside $\mathcal{C}_R^{ass}$ by $\mathcal{C}_{R/A}^{ass}$. The category $\mathcal{C}_{R/A}^{ass}$ has a topological model category structure inherited from $\mathcal{C}_R^{ass}$.

Let M be an A -bimodule. Then we could form the R -algebra with the underlying R -module $A \vee M$, the ‘square-zero extension’ of A by M . There is an obvious product and augmentation maps making $A \vee M$ into an R -algebra over A .

We now introduce the analogue of the abelianization functor in the noncommutative context. Let B be a cofibrant R -algebra over A and denote by $\Omega_A^{ass}(B)$ the A -bimodule

$$A \wedge_B \Omega_B^{ass} \wedge_B A \cong A \wedge A^{op} \wedge_{B \wedge B^{op}} \Omega_B^{ass}.$$

Notice that $\Omega_A^{ass}(B)$ is an A -bimodule.

Theorem 3.1. *There is a natural equivalence*

$$h\mathcal{C}_{R/A}^{ass}(B, A \vee M) \cong h\mathcal{M}_{A-A}(\Omega_A^{ass}(B), M).$$

Let us show how to associate to any map $B \longrightarrow A \vee M$ in $ho\mathcal{C}_{R/A}^{ass}$ a map of A -bimodules $\Omega_A^{ass}(B) \longrightarrow M$.

Denote by $\overline{A \vee M}$ the fibrant cofibrant approximation of $A \vee M$ in the category of R -algebras over A . It suffices to construct a ‘universal’ map of $\overline{A \vee M}$ -bimodules

$$(3.2) \quad \Omega_A^{ass}(\overline{A \vee M}) \longrightarrow M.$$

Indeed the for a map $B \longrightarrow \overline{A \vee M}$ in $\mathcal{C}_{R/A}^{ass}$ we have a composite map

$$\Omega_A^{ass}(B) \longrightarrow \Omega_A^{ass}(\overline{A \vee M}) \longrightarrow M$$

and therefore a correspondence

$$(3.3) \quad h\mathcal{C}_{R/A}^{ass}(B, \overline{A \vee M}) = h\mathcal{C}_{R/A}^{ass}(B, A \vee M) \longrightarrow h\mathcal{M}_{A-A}(\Omega_A^{ass}(B), M)$$

as desired.

Further since

$$\Omega_A^{ass}(\overline{A \vee M}) = A \wedge_{\overline{A \vee M}} \Omega_{\overline{A \vee M}}^{ass} \wedge_{\overline{A \vee M}} A$$

a map of (3.2) is the same as a map of $\overline{A \vee M}$ -bimodules $\Omega_{\overline{A \vee M}}^{ass} \longrightarrow M$. Instead of the latter map we construct a map $\Omega_{A \vee M}^{ass} \longrightarrow M$. This would be good enough since even $A \vee M$ is not a cofibrant object in $\mathcal{C}_{R/A}^{ass}$ the smash product $(A \vee M) \wedge (A \vee M)$ clearly represents the derived smash product and therefore $\Omega_{\overline{A \vee M}}^{ass}$ is weakly equivalent to $\Omega_{A \vee M}^{ass}$. Consider the following diagram in the homotopy category of $A \vee M$ -bimodules:

$$\begin{array}{ccc} A \wedge A \vee A \wedge M \vee M \wedge A \vee M \wedge M & \longrightarrow & A \vee M \\ \downarrow & & \parallel \\ A \vee M \vee M & \longrightarrow & A \vee M \end{array}$$

Here the left vertical arrow is determined by the R -algebra structure on A , an A -bimodule structure on M and is zero on the last summand. The lower horizontal arrow is zero on A , identity on the first M -summand and minus identity on the last M -summand. Then the homotopy fibre of the upper row is equivalent to $\Omega_{A \vee M}^{ass}$ by definition and the homotopy fibre of the lower row is equivalent to M . There results a map of $A \vee M$ -bimodules $\Omega_{A \vee M}^{ass} \longrightarrow M$ as desired.

The proof of the theorem is then completed by showing that the above correspondence (3.3) is in fact an equivalence if B is equal to the free algebra

$$B = TV = R \vee V \vee V^{\otimes 2} \vee \dots$$

where V is a cofibrant R -module over A and then resolving a general B by a monadic bar-construction. Note that the homotopy fibre of the multiplication map $TV \wedge TV \longrightarrow TV$ is equivalent to the TV -bimodule $TV \wedge V \wedge TV$ and therefore

$$\Omega_{TV}^{ass} \simeq TV \wedge V \wedge TV.$$

We have the ‘universal derivation’ map $A \longrightarrow A \vee \Omega_A^{ass}$ which is adjoint to the identity map $\Omega_A^{ass}(A) = \Omega_A^{ass} \longrightarrow \Omega_A^{ass}$. Recall that as an R -module Ω_A^{ass} (even as a left A -module) is weakly equivalent to $A \wedge A/R$. Taking the composition of the universal derivation with the canonical projection $A \vee \Omega_A^{ass} \longrightarrow \Omega_A^{ass}$ we get a map $\phi : A \longrightarrow \Omega_A^{ass} \simeq A \wedge A/R$. Then as a map of R -modules ϕ is homotopic to the map

$$A \cong R \wedge A \xrightarrow{1 \wedge d} A \wedge A/R$$

where d is defined as the second arrow in the homotopy cofibre sequence

$$R \xrightarrow{1} A \xrightarrow{d} A/R.$$

This could be seen by first taking A to be the free algebra TV and then resolving a general R -algebra A by a simplicial construction consisting of free R -algebras.

Let us denote by $Der_R^0(A, M)$ the set of homotopy classes of maps $A \longrightarrow A \vee M$ in $h\mathcal{C}_{A/R}^{ass}$. We will call elements of $Der_R^0(A, M)$ *topological derivations* of A with values in the A -bimodule M . Then Theorem 3.1 shows that $Der_R^0(A, M)$ is in fact an abelian group. Moreover, $Der_R^0(A, M)$ is the zeroth homotopy group of the function spectrum $F_{A \wedge A^{op}}(\Omega_A^{ass}, M)$. We will call it the spectrum of topological derivations of A with values in M and denote it by $\mathbf{Der}_R(A, M)$. The homotopy groups of $\mathbf{Der}_R(A, M)$ will be denoted by $Der_R^i(A, M)$, so $Der_R^i(A, M) = \pi_{-i} \mathbf{Der}_R(A, M)$.

Remark 3.2. Arguing in the same way as for commutative derivations we see that there is a weak equivalence of topological spaces

$$\mathrm{Map}_{\mathcal{C}_{R/A}^{ass}}(A, A \vee M) \simeq \Omega^\infty \mathbf{Der}_R(A, M).$$

In particular the space $\mathrm{Map}_{\mathcal{C}_{R/A}^{ass}}(A, A \vee M)$ is an infinite loop space.

Let $\tilde{A}$ be a cofibrant replacement of A as an A -bimodule. Replacing A with $\tilde{A}$ in the homotopy fibre sequence (3.1) and applying the functor $F_{A \wedge A^{op}}(?, M)$ to it we obtain the homotopy fibre sequence:

$$F_{A \wedge A^{op}}(\tilde{A}, M) \longrightarrow M \longrightarrow \mathbf{Der}_R(A, M).$$

The function spectrum $F_{A \wedge A^{op}}(\tilde{A}, M)$ is called *topological Hochschild cohomology* spectrum of A with values in M and has a special notation $\mathbf{THH}_R(A, M)$. So we obtain the following homotopy fibre sequence relating topological Hochschild cohomology and topological derivations:

$$(3.4) \quad \mathbf{THH}_R(A, M) \longrightarrow M \longrightarrow \mathbf{Der}_R(A, M).$$

This fibre sequence to a large extent reduces the calculation of $Der_R^*(A, M)$ to that of topological Hochschild cohomology groups $THH_R^*(A, M) := \pi_{-i} \mathbf{THH}_R(A, M)$. The latter groups are relatively computable due to the hypercohomology spectral sequence of [7]:

$$Ext_{(A \wedge A^{op})_*}^{**}(A_*, M_*) \implies THH_R^*(A, M).$$

Similarly to the commutative case there is a forgetful map

$$l_R^* : Der_R^*(A, M) \longrightarrow [A, M]_{R-mod}^*$$

whose image is contained in the subspace of primitive operations in $[A, M]_{R-mod}^*$.

Remark 3.3. There is also the notion of topological Hochschild *homology* spectrum

$$\mathbf{THH}^R(A, M) := \tilde{A} \wedge_{A \wedge A^{op}} M.$$

Topological Hochschild homology plays an important role in computations of algebraic K -theory of rings, cf. [20]. If the R -algebra A is commutative then both $\mathbf{THH}_R$ and $\mathbf{THH}^R$ are R -modules and there is a duality between them in $h\mathcal{M}_A$:

$$\mathbf{THH}_R(A, M) \simeq F_A(\mathbf{THH}^R(A, M), A).$$

Remark 3.4. Just as topological André-Quillen cohomology the spectrum $\mathbf{Der}_R(A, M)$ is covariant with respect to M and contravariant with respect to A . The arguments are precisely the same as in the commutative context except that we refer to Theorem 3.1 instead of Theorem 2.4 and A -modules are replaced with A -bimodules. For a map of A -bimodules $f : M \longrightarrow N$ and a topological derivation $d : A \longrightarrow A \vee M$ we have a topological derivation $f_* d$ of A with values in N . For a map of R -algebras $g : X \longrightarrow A$ we obtain a topological derivation $g^* d$ of X with values in M .

Sometimes in the noncommutative context it is useful to consider the so-called *relative topological derivations* which we will now define. Let R' be a not necessarily commutative R -algebra. Denote by $\mathcal{C}_{R'}^{ass}$ the category of R -algebras A supplied with a fixed R -algebra map $R' \longrightarrow A$, not necessarily central. In other words $\mathcal{C}_{R'}^{ass}$ is the undercategory of R' in $\mathcal{C}_R$. The objects of $\mathcal{C}_{R'}^{ass}$ will be called *R' -algebras* by a slight abuse of language. Let $m : A \wedge A \longrightarrow A$ be the multiplication map. Note that there exists a unique map $m' : A \wedge_{R'} A \longrightarrow A$ such that the diagram

$$\begin{array}{ccc} A \wedge A & \xrightarrow{m} & A \\ & \searrow & \nearrow m' \\ & A \wedge_{R'} A & \end{array}$$

is commutative.

Now suppose without loss of generality that the R -algebra R' is cofibrant and that the structure map $R' \longrightarrow A$ is a cofibration of R -algebras. We define the R' -relative module of differentials $\Omega_{A/R'}^{ass}$ from the following homotopy fibre sequence:

$$\Omega_{A/R'}^{ass} \longrightarrow A \wedge_{R'} A \xrightarrow{m'} A.$$

Let $\mathcal{C}_{A/R'}^{ass}$ denote the category whose objects are R' -algebras B supplied with an R' -algebra map $B \longrightarrow A$. Note that if M is an A -bimodule then $A \vee M$ is an object of $\mathcal{C}_{A/R'}^{ass}$. This gives a functor $\mathcal{M}_{A-A} \longrightarrow \mathcal{C}_{A/R'}^{ass}$. It turns out that on the level of homotopy categories this functor admits a left adjoint

$$B \longrightarrow \Omega_{A/R'}^{ass}(B) := A \wedge_B \Omega_{B/R'}^{ass} \wedge_B A.$$

More precisely there is a natural equivalence

$$h\mathcal{C}_{A/R'}^{ass}(B, A \vee M) \cong h\mathcal{M}_{A-A}(\Omega_{A/R'}^{ass}(B), M).$$

The proof is similar to the one in the absolute case. One makes use of the ‘free’ R' -algebra functor $T_{R'}$ given for an R' -bimodule V as

$$T_{R'}V = R' \vee (V \vee V \wedge_{R'} V) \vee \dots$$

instead of the usual free R -algebra on an R -module V .

The homotopy classes of maps in $h\mathcal{C}_{A/R'}^{ass}(A, A \vee M)$ are called topological derivations of A with values in M *relative to* R' . An algebraic analogue of relative topological derivations are the derivations of an algebra A with coefficients in an A -bimodule M vanishing on a subalgebra R' of A . Relative modules of differentials were studied in the context of noncommutative differential forms in [5]. Replacing $\Omega_{R/R'}^{ass}$ with $\tilde{\Omega}_{R/R'}^{ass}$, its cofibrant approximation as an A -bimodule we define the R -module of R' -relative derivations of A with values in an A -bimodule M :

$$\mathbf{Der}_{R/R'}(A, M) := F_{A \wedge A^{op}}(\tilde{\Omega}_{R/R'}^{ass}, M).$$

It is easy to see that there is the following homotopy fibre sequence of R -modules:

$$\mathbf{THH}_R(A, M) \longrightarrow \mathbf{THH}_R(R', M) \longrightarrow \mathbf{Der}_{R/R'}(A, M)$$

which specializes to (3.4) when $R = R'$.

4. OBSTRUCTION THEORY

In this section we show that topological Andre-Quillen cohomology in the commutative case and topological derivations in the associative case are a natural home for lifting algebra maps. For definiteness we work in the context of associative algebras, however the results have obvious analogues, with the same proofs, in the commutative case. Topological derivations are replaced with commutative derivations, spectra $\mathbf{Der}_R(-, -)$ - with spectra $\mathbf{TAQ}_R(-, -)$ and the spaces $\mathrm{Map}_{\mathcal{C}_R^{ass}}(-, -)$ - with spaces $\mathrm{Map}_{\mathcal{C}_R^{comm}}(-, -)$.

Let A be a cofibrant R -algebra, I be an A -bimodule and $d : A \longrightarrow A \vee \Sigma I$ be a derivation of A with coefficients in ΣI , the suspension of I . Let $\epsilon : A \longrightarrow A \vee \Sigma A$ be the inclusion of a retract. Define the R -algebra B from the following homotopy pullback square of R -algebras:

$$\begin{array}{ccc} B & \longrightarrow & A \\ \downarrow & & \downarrow \epsilon \\ A & \xrightarrow{d} & A \vee \Sigma I \end{array}$$

It is clear that the homotopy fibre of the map A is weakly equivalent to the A -bimodule I .

Definition 4.1. The homotopy fibre sequence

$$I \longrightarrow B \longrightarrow A$$

is called the topological singular extension of A by I associated with the derivation $d : A \longrightarrow A \vee \Sigma I$.

Theorem 4.2. *Let $I \rightarrow B \rightarrow A$ be a singular extension of R -algebras associated with a derivation $d : B \rightarrow B \vee \Sigma I$ and $f : X \rightarrow A$ a map of R -algebras where the R -algebra X is cofibrant. Then:*

- (1) *The map f lifts to an R -algebra map $X \rightarrow B$ if and only if the induced derivation $f^*d \in \mathrm{Der}_R^1(X, I) = \mathrm{Der}_R^0(X, \Sigma I)$ is homotopic to zero.*
- (2) *Assuming that a lifting exists in the fibration*

$$(4.1) \quad \mathrm{Map}_{\mathcal{C}_R^{ass}}(X, B) \rightarrow \mathrm{Map}_{\mathcal{C}_R^{ass}}(X, A)$$

the homotopy fibre over the point $f \in \mathrm{Map}_{\mathcal{C}_R^{ass}}(X, A)$ is weakly equivalent to $\Omega^\infty \mathbf{Der}_R(X, I)$ (the 0th space of the spectrum $\mathbf{Der}_R(X, I)$).

The proof is rather formal. Assume first that a lifting of f exists. Then $d \circ f$ factors as

$$X \xrightarrow{f} B \longrightarrow A \xrightarrow{\epsilon} A \vee \Sigma I$$

which means that the derivation $d \circ f$ is trivial. Conversely, if $d \circ f$ is trivial then the diagram of R -algebras

$$\begin{array}{ccc} X & \xrightarrow{f} & A \\ \downarrow f & & \downarrow \epsilon \\ A & \xrightarrow{d} & A \vee \Sigma I \end{array}$$

commutes up to homotopy and by the universal property of the homotopy pullback there is a map $\tilde{f} : X \rightarrow B$ lifting f .

For the second part suppose that the lifting of f exists, so the (homotopy) fibre of the map (4.1) is nonempty. We have the following diagram of R -algebras.

$$\begin{array}{ccc} B & \xrightarrow{\quad} & A \\ & \searrow & \nearrow \\ & A & \\ & \nearrow & \searrow \\ A & \xrightarrow{d} & A \vee \Sigma I \end{array}$$

Changing B in its homotopy class if necessary we can arrange that this diagram be strictly commutative. Notice that the outer square is a homotopy pullback of R -algebras. Applying the functor $\text{Map}_{\mathcal{C}_R^{ass}}(X, -)$ to this diagram we get the diagram of spaces

$$\begin{array}{ccccc} \text{Map}_{\mathcal{C}_R^{ass}}(X, B) & \xrightarrow{\quad} & \text{Map}_{\mathcal{C}_R^{ass}}(X, A) & & \\ & \searrow & \nearrow & & \\ & \text{Map}_{\mathcal{C}_R^{ass}}(X, A) & & & \\ & \nearrow & \searrow & & \\ \text{Map}_{\mathcal{C}_R^{ass}}(X, A) & \xrightarrow{\quad} & \text{Map}_{\mathcal{C}_R^{ass}}(X, A \vee \Sigma I) & & \end{array}$$

Again the outer square is a homotopy pullback (of spaces). Taking the homotopy fibres of the maps from the outer square to the center (over $f \in \text{Map}_{\mathcal{C}_R^{ass}}$) we get the following homotopy pullback of spaces:

$$(4.2) \quad \begin{array}{ccccc} \text{hofib Map}_{\mathcal{C}_R^{ass}}(X, B) & \rightarrow & \text{Map}_{\mathcal{C}_R^{ass}}(X, A) & \xrightarrow{\quad} & pt \\ \downarrow & & & & \downarrow \\ pt & \xrightarrow{\quad} & \text{hofib Map}_{\mathcal{C}_R^{ass}}(X, A \vee \Sigma I) & \rightarrow & \text{Map}_{\mathcal{C}_R^{ass}}(X, A) \end{array}$$

Notice that the space in the right lower corner of (4.2) is canonically weakly equivalent to $\Omega^\infty \mathbf{Der}_R(X, \Sigma I)$, see Remark 3.2. Since according to our assumption the space

$$\text{hofib Map}_{\mathcal{C}_R^{ass}}(X, B) \longrightarrow \text{Map}_{\mathcal{C}_R^{ass}}(X, A)$$

is nonempty the images of the lower and right arrows in (4.2) coincide and therefore

$$\begin{aligned} \mathrm{hofib} \, \mathrm{Map}_{C_R^{\mathrm{ass}}}(X, B) &\rightarrow \mathrm{Map}_{C_R^{\mathrm{ass}}}(X, A) \simeq \Omega(\Omega^\infty \mathbf{Der}_R(X, \Sigma I)) \\ &\simeq \Omega^\infty \mathbf{Der}_R(X, I). \end{aligned}$$

This finishes the proof of Theorem 4.2.

A large supply of topological singular extensions can be obtain by taking Postnikov stages of connective R -algebras of commutative R -algebras. We have the following

Theorem 4.3. *Let A be a connective R -algebra where R is a connective commutative S -algebra. Then there exists a tower of R -algebras under A :*

$$\begin{array}{ccccccc} & & A & & & & \\ & \swarrow f_0 & & \searrow f_{n+1} & & & \\ A_0 & \longleftarrow A_1 & \longleftarrow \cdots & \longleftarrow A_n & \longleftarrow A_{n+1} & \longleftarrow \cdots \end{array}$$

such that

- $\pi_k A_n = 0$ for $k > n$;
- the map $f_n : A \rightarrow A_n$ induces an isomorphism $\pi_k A \rightarrow \pi_k A_n$ for $0 \leq k \leq n$;
- the homotopy fibre sequences $H\pi_{n+1} A \rightarrow A_{n+1} \rightarrow A_n$ are topological singular extensions associated with topological derivations $k_n \in \mathrm{Der}_R^1(A_n, H\pi_{n+1} A)$.

In the case when A is a connective commutative R -algebra we have the precise analogue of the theorem above: there is a tower $\{A_n\}$ of *commutative* R -algebras under A and the homotopy fibre sequences $H\pi_{n+1} A \rightarrow A_{n+1} \rightarrow A_n$ are topological singular extensions associated with *commutative* derivations $k_n \in \mathrm{TAQ}_R^1(A_n, H\pi_{n+1} A)$.

Theorem 4.3 is proved by induction on n . By glueing cells in the category of R -algebras or commutative R -algebras we construct a map $A \rightarrow A_0 = H\pi_0 A$. Suppose that the n th stage A_n for a commutative R -algebra A_n is already constructed. It is not hard to see using the spectral sequence (2.1) that the lowest nonzero homotopy group of $\Omega_{A_n/A}^{\mathrm{comm}}$ is in dimension $n+1$ and is equal to $\pi_{n+1} A$. We then construct a map $k_n : \Omega_{A_n/A}^{\mathrm{comm}} \rightarrow H\pi_{n+1} A$ by attaching cells to $\Omega_{A_n/A}^{\mathrm{comm}}$ to kill its higher homotopy groups. The map k_n is the required commutative derivation. In the associative context we use the A -bimodule of relative differentials $\Omega_{A_n/A}^{\mathrm{ass}}$ and proceed similarly.

Theorem 4.3 provides a large supply of R -algebras or commutative R -algebras. For example Postnikov stages of MU or kU are commutative S -algebras. Other examples of commutative S -algebras include the Morava E -theory spectra E_n and Eilenberg-MacLane spectra Hk for a commutative ring k . It is generally very hard to prove that a spectrum possesses a commutative S -algebra structure unless there is a geometric reason for it. We will see later on that the situation with associative structures is better; a large class of complex-oriented spectra could be given structures of MU -algebras and, therefore, S -algebras.

5. STABILIZATION

In this section we give an interpretation of topological derivation and topological André-Quillen cohomology as ‘stabilizations’ of a certain forgetful functors. This interpretation in the commutative case is due to Basterra-McCarthy [2] and Basterra-Mandell (private communication).

Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a continuous functor between pointed topological model categories. We assume that F is a homotopy functor which means that it preserves homotopy equivalences. We also assume that F is reduced, that is it takes the initial object of $\mathcal{C}$ into a contractible object in $\mathcal{D}$ (i.e. homotopy equivalent to the initial object in $\mathcal{D}$). For an object X in $\mathcal{D}$ we will denote by ΩX the cotensor of X and the pointed circle S^1 . In addition the tensor of $?$ with a pointed topological space X will be denoted by $X \hat{\otimes} ?$.

We have the following pushout diagram of pointed spaces:

$$(5.1) \quad \begin{array}{ccc} S^0 & \longrightarrow & I \\ \downarrow & & \downarrow \\ I & \longrightarrow & S^1 \end{array}$$

where I is the unit interval. Tensoring it with X and applying the functor F we get the diagram

$$\begin{array}{ccc} F(X) = F(S^0 \hat{\otimes} X) & \longrightarrow & F(I \hat{\otimes} X) \\ \downarrow & & \downarrow \\ F(I \hat{\otimes} X) & \longrightarrow & F(S^1 \hat{\otimes} X) \end{array}$$

Define the functor $X \longrightarrow TF(X)$ by the requirement that the diagram

$$\begin{array}{ccc} TF(X) & \longrightarrow & F(I \hat{\otimes} X) \\ \downarrow & & \downarrow \\ F(I \hat{\otimes} X) & \longrightarrow & F(S^1 \hat{\otimes} X) \end{array}$$

be a homotopy pullback in $\mathcal{D}$. There results a natural transformation $\xi : F(X) \longrightarrow TF(X)$.

Definition 5.1. The stabilization of F is the functor F^{st} defined as

$$F^{st}(X) = \text{hocolim}(F(X) \longrightarrow TF(X) \longrightarrow TTF(X) \longrightarrow \dots)$$

where the colimit is taken over the maps $T^i(\xi) : T^i(F(X)) \longrightarrow T^i(TF(X))$.

Note that since $I \hat{\otimes} X$ is contractible the object $TF(X)$ is homotopy equivalent to $\Omega F(S^1 \hat{\otimes} X)$. Furthermore, it is easy to see that there is a homotopy equivalence

$$T^n FX \simeq \Omega^n F(S^n \hat{\otimes} X)$$

where Ω^n is the n th iterate of Ω .

Remark 5.2. The stabilization (or linearization) of a homotopy functor was introduced by Goodwillie in [11] and constitutes the first layer of its *Taylor tower*, [12]. It would be interesting to investigate the higher layers of this tower in our context.

We are now ready to give the interpretation of Der and TAQ as stabilizations of certain forgetful functors. We start with the associative case. Let A be a cofibrant R -algebra and consider the category $\mathcal{C}_{A/A}^{ass}$ of A -algebras over A . Recall that $\mathcal{C}_{A/A}^{ass}$ consists of R -algebras B supplied with two maps $\eta_B : B \longrightarrow A$ and $\zeta_B : A \longrightarrow B$ such that $\eta_A \circ \zeta_A = id_A$. The category $\mathcal{C}_{A/A}^{ass}$ is a topological model category. Let us first describe its cotensors. Let $\tilde{B}$ be a fibrant approximation of B as an object in $\mathcal{C}_{A/A}^{ass}$. Then $\eta_{\tilde{B}} : \tilde{B} \longrightarrow A$ is a fibration of R -algebras. Denote by $I(B)$ the fibre of $\eta_{\tilde{B}}$ so that there is a homotopy fibre sequence of B -bimodules:

$$I(B) \longrightarrow B \xrightarrow{\eta_B} A.$$

Consider the R -module $A \vee I(B)$. It is clearly an object in $\mathcal{C}_{A/A}^{ass}$ and, moreover, the obvious map $A \vee I(B) \longrightarrow B$ is a weak equivalence. Then for a based space X the R -algebra B^X , the cotensor of B and X in $\mathcal{C}_{A/A}^{ass}$ is weakly equivalent to $A \vee I(B)^X$ where $I(B)^X = F_S(\Sigma^\infty X, I(B))$ is the usual cotensor (function spectrum) of X and the R -module $I(B)$.

Now we will describe tensors in $\mathcal{C}_{A/A}^{ass}$. Let B be a cofibrant object in $\mathcal{C}_{A/A}^{ass}$ and X be a based space that consists of $n + 1$ points, one of them being the base point. Then the tensor $X \hat{\otimes} B$

is just the n -fold free product of B over A : $B \amalg_A^n$. For a general based space X we replace it with the simplicial set X_* whose realization is equivalent to X , and tensor it component-wise with B . The realization of the obtained simplicial object in $\mathcal{C}_{A/A}^{ass}$ will be homotopy equivalent to $X \hat{\otimes} B$. For example if we take $X = S^1$, the circle and by X_* its simplicial model having two nondegenerate simplices in dimensions 0 and 1 then the resulting simplicial object $X_* \hat{\otimes} B$ is just the bar-construction $\beta_*(A, B, A)$. We have

$$\beta_i(A, B, A) = B \amalg_A^i.$$

The face maps are induced in the usual way by the canonical folding map $B \amalg_A B \rightarrow B$ and the degeneracies are induced by the map ζ_B .

Recall that the functor $\Omega_{B/A}^{ass}$ is defined from the fibre sequence

$$\Omega_{B/A}^{ass} \longrightarrow B \wedge_A B \xrightarrow{m'} B.$$

We will assume that $\Omega_{B/A}^{ass}$ is a cofibrant B -bimodule or else replace it by its cofibrant approximation. By definition,

$$\Omega_{A/A}^{ass}(B) := A \wedge_B \Omega_{B/A}^{ass} \wedge_B A.$$

Both I and $\Omega_{A/A}^{ass}(B)$ considered as functors of B take their values in the category of A -bimodules.

Theorem 5.3. *Let A and B be connective R -algebras. Then $I^{st}(B) \simeq \Omega_{A/A}^{ass}(B)$*

To see this consider the ‘universal derivation’ map $B \rightarrow A \vee \Omega_{A/A}^{ass}(B)$ and the composition

$$f_B : I(B) \longrightarrow B \longrightarrow A \vee \Omega_{A/A}^{ass}(B) \longrightarrow \Omega_{A/A}^{ass}(B).$$

It is easy to see that there is a weak equivalence of R -modules

$$\Omega_{A/A}^{ass}(B) \simeq A \wedge_B I(B)$$

and the map f_B could be represented as

$$I(B) \cong B \wedge_B I(B) \xrightarrow{\eta_B \wedge id} A \wedge_B I(B).$$

Note that if the $\eta_B : B \rightarrow A$ is n -connected then the map f_B is $2n + 1$ -connected. Further if $I(B)$ is m -connected then $I(S^n \hat{\otimes} B)$ is $n + m$ -connected and the map

$$f_{S^n \hat{\otimes} B} : I(S^n \hat{\otimes} B) \longrightarrow \Omega_{A/A}^{ass}(S^n \hat{\otimes} B)$$

is $2(m + n) + 1$ -connected. It follows that the map

$$\Omega^n f_{S^n \hat{\otimes} B} : \Omega^n I(S^n \hat{\otimes} B) \longrightarrow \Omega^n \Omega_{A/A}^{ass}(S^n \hat{\otimes} B)$$

is $2m + 1$ -connected and therefore $I^{st}(B) = [\Omega_{A/A}^{ass}]^{st}(B)$. The results of Section 2 imply that the functor $B \rightarrow [\Omega_{A/A}^{ass}]^{st}(B)$ preserves tensors and therefore

$$\Omega^n \Omega_{A/A}^{ass}(S^n \hat{\otimes} B) \simeq \Omega^n \Sigma^n \Omega_{A/A}^{ass}(B) \simeq \Omega_{A/A}^{ass}(B),$$

which implies that $[\Omega_{A/A}^{ass}]^{st}(B) \simeq \Omega_{A/A}^{ass}(B)$. This finishes our sketch of the proof of Theorem 5.3.

What is the relation of the functor $\Omega_{A/A}^{ass}(B)$ to the topological derivations of A ? Let $B = A \amalg A$.

Define ζ_B to be the map $A = A \amalg R \xrightarrow{id \amalg 1} A \amalg A$ and $\eta_B : A \amalg A \rightarrow A$ to be the folding map. Then B becomes an object in $\mathcal{C}_{A/A}^{ass}$. Moreover we have a natural isomorphism for any $C \in \mathcal{C}_{A/A}^{ass}$:

$$h\mathcal{C}_{A/A}^{ass}(B, C) \cong h\mathcal{C}_{R/A}(A, C).$$

Therefore for an A -bimodule M we have

$$\begin{aligned} \mathrm{Der}_R^0(A, M) &= h\mathcal{C}_{R/A}(A, A \vee M) \cong h\mathcal{C}_{A/A}^{ass}(B, A \vee M) \\ &\cong h\mathcal{M}_{A-A}(\Omega_{A/A}^{ass}, M) \\ &\cong h\mathcal{M}_{A-A}(I^{st}(B), M). \end{aligned}$$

We can also obtain in the usual way the enriched version of the above equivalence:

$$\mathbf{Der}_R(A, A \vee M) \simeq F_{A \wedge A^{op}}(I^{st}(A \coprod A), M).$$

Theorem 5.3 has an analogue in the context of commutative R -algebras. The corresponding result was first established by M.Basterra and M.Mandell, cf, also [2] for TAQ and our proof was modeled on theirs. One should also mention that the original definition of TAQ due to I. Kriz [15] was in terms of stabilization. In contrast with the associative case this interpretation is extremely helpful in concrete calculations.

Let us now describe the result of Basterra and Mandell. A is now a commutative cofibrant R -algebra. Consider the category $\mathcal{C}_{A/A}^{comm}$ of commutative A -algebras over A . Then $I(B) \in \mathcal{M}_A$ is defined from the fibre sequence

$$I(B) \longrightarrow \tilde{B} \xrightarrow{\eta_{\tilde{B}}} A$$

where $\tilde{B}$ is the fibrant replacement of B in $\mathcal{C}_{A/A}^{comm}$. Recall the definition of the *indecomposables functor* $B \rightarrow Q(B)$ from Section 1. It turns out that if A and B are connected then $I^{st}(B)$, the stabilization of the functor I evaluated at B , is weakly equivalent to $Q \circ I(B)$. The proof is very similar to the one given in the associative case. To show that the map

$$I(S^n \hat{\otimes} B) \longrightarrow Q \circ I(S^n \hat{\otimes} B)$$

is sufficiently highly connected one uses Proposition 2.2. To relate stabilization with TAQ consider $A \wedge A$ as an A -algebra over A . The augmentation $A \wedge A \rightarrow A$ is just the multiplication map and the A -algebra structure on $A \wedge A$ is defined via the action of A on the left smash factor. Then for an A -module M we have the following isomorphisms in $h\mathcal{M}_A$:

$$\mathbf{TAQ}_R(A, M) \simeq F_A(Q \circ I(A \wedge A), M) \simeq F_A(I^{st}(A \wedge A), M).$$

In fact the author was informed by M.Basterra that the last isomorphism holds even without the hypothesis that A is connective. That suggests that Theorem 5.3 might also hold without any connectivity assumptions.

6. CALCULATIONS

Up until now we considered only formal properties of TAQ and topological derivations and saw that they are quite similar. The difference appears in their calculational aspect. Topological André-Qullen cohomology is far harder to compute and presently very few explicit calculations have been carried out. By contrast, topological derivations are relatively approachable. In most cases their computation reduces to that of topological Hochschild cohomology thanks to the homotopy fibre sequence (3.4). There is an extensive literature dedicated to the computation of THH of different Eilenberg-MacLane spectra of which we mention [4], [8], [19]. Topological Hochschild (co)homology of some spectra other than Eilenberg-MacLane are computed in [24] and [3]. We do not intend to give a detailed review of this literature and restrict ourselves with giving a few examples related to obstruction theory. In this section the symbol ' $\wedge$ ' stands for ' $\wedge_S$ '.

Our first example is the computation of topological derivations of $H\mathbb{F}_p$.

Theorem 6.1. (1) *There is an isomorphism of graded $\mathbb{F}_p$ -vector spaces*

$$THH_S^*(\mathbb{F}_p, \mathbb{F}_p) \cong (\mathbb{F}_p[x])^* = \text{Hom}_{\mathbb{F}_p}(\mathbb{F}_p[x], \mathbb{F}_p).$$

where x has degree 2.

(2) *There is an isomorphism of graded $\mathbb{F}_p$ -vector spaces*

$$\text{Der}_S^{*-1}(\mathbb{F}_p, \mathbb{F}_p) \cong (\mathbb{F}_p[x])^*/(\mathbb{F}_p)$$

Note that the statement (2) of the theorem follows from statement (1) because the homotopy fibre sequence (3.4) in our case splits giving the short exact sequence

$$\mathbb{F}_p \longrightarrow THH_S^*(\mathbb{F}_p, \mathbb{F}_p) \longrightarrow \text{Der}_S^{*-1}(\mathbb{F}_p, \mathbb{F}_p).$$

To obtain the claim about $THH_S^*(\mathbb{F}_p, \mathbb{F}_p)$ it is enough to compute topological Hochschild *homology* $THH_*^S(\mathbb{F}_p, \mathbb{F}_p)$ because

$$THH_S^*(\mathbb{F}_p, \mathbb{F}_p) \cong \text{Hom}_{\mathbb{F}_p}(THH_*^S(\mathbb{F}_p, \mathbb{F}_p), \mathbb{F}_p),$$

see Remark 3.3. The computation of $THH_*^S(\mathbb{F}_p, \mathbb{F}_p)$ is due to Bökstedt, [4]. Briefly, one uses the spectral sequence

$$E_{**}^2 = \text{Tor}_{H\mathbb{F}_p \wedge H\mathbb{F}_p}(\mathbb{F}_p, \mathbb{F}_p) = \text{Tor}_{\mathcal{A}_*^p}(\mathbb{F}_p, \mathbb{F}_p) \Rightarrow THH_*^S(\mathbb{F}_p, \mathbb{F}_p)$$

where $\mathcal{A}_*^p = H\mathbb{F}_p \wedge H\mathbb{F}_p$ is the dual Steenrod algebra mod p . The E_2 -term of this spectral sequence is easy to compute. It turns out that for $p = 2$ there are no further differentials and for an odd p one uses the Dyer-Lashof operations to compute the only nontrivial differential d^{p-1} . The same operations also solve the extension problem. The final result is that, regardless of p ,

$$(6.1) \quad THH_*^S(H\mathbb{F}_p, H\mathbb{F}_p) \cong \mathbb{F}_p[x]$$

with x in homological degree 2. This finishes our sketch of Theorem 6.1.

Remark 6.2. In fact, Bökstedt's calculation shows that the isomorphism of (6.1) is multiplicative. This fact will be used in the computation of $TAQ_S^*(H\mathbb{F}_p, H\mathbb{F}_p)$.

Our next example is the calculation of $\text{Der}_{MU}^*(H\mathbb{F}_p, H\mathbb{F}_p)$ where MU is the complex cobordism spectrum. Recall that the coefficient ring of MU is the polynomial algebra:

$$MU_* = \mathbb{Z}[x_1, x_2, \dots]$$

where $|x_i| = 2i$. Since MU is known to be a commutative connective S -algebra the spectrum $H\mathbb{F}_p$ is naturally an MU -algebra and it makes sense to consider topological derivations and topological Hochschild cohomology of $H\mathbb{F}_p$ as an MU -algebra. The relevant result is

Theorem 6.3. (1) *There is an isomorphism of graded $\mathbb{F}_p$ -vector spaces*

$$THH_{MU}^*(H\mathbb{F}_p, H\mathbb{F}_p) = \mathbb{F}_p[y_1, y_2, \dots]$$

where y_i has cohomological degree $2i$.

(2) *There is an isomorphism of graded $\mathbb{F}_p$ -vector spaces*

$$\text{Der}_{MU}^{*-1}(H\mathbb{F}_p, H\mathbb{F}_p) = \mathbb{F}_p[y_1, y_2, \dots]/(\mathbb{F}_p).$$

Again the second claim is a consequence of the first one. To verify the first claim we need to compute $\pi_* H\mathbb{F}_p \wedge_{MU} H\mathbb{F}_p$. Using the spectral sequence

$$\begin{aligned} \text{Tor}_{**}^{MU}(\mathbb{F}_p, \mathbb{F}_p) &= \text{Tor}_{**}^{\mathbb{Z}[x_1, x_2, \dots]}(\mathbb{F}_p, \mathbb{F}_p) \\ &\Rightarrow \pi_* H\mathbb{F}_p \wedge_{MU} H\mathbb{F}_p \end{aligned}$$

we see that the ring $\pi_* H\mathbb{F}_p \wedge_{MU} H\mathbb{F}_p$ is isomorphic to $\Lambda_{\mathbb{F}_p}(z_1, z_2, \dots)$, the exterior algebra over $\mathbb{F}_p$ on generators z_i where $|z_i| = 2i - 1$.

Finally from the spectral sequence

$$\begin{aligned} Ext_{\pi_* H\mathbb{F}_p \wedge_{MU} H\mathbb{F}_p}^{**}(\mathbb{F}_p, \mathbb{F}_p) &= Ext_{\Lambda_{\mathbb{F}_p}^{**}(z_1, z_2, \dots)}^{**}(\mathbb{F}_p, \mathbb{F}_p) \\ &= \mathbb{F}_p[y_1, y_2, \dots] \Rightarrow THH_{MU}^*(H\mathbb{F}_p, H\mathbb{F}_p) \end{aligned}$$

we obtain the desired isomorphism.

Next we would like to discuss the forgetful maps

$$l_{MU}^* : Der_{MU}^*(H\mathbb{F}_p, H\mathbb{F}_p) \longrightarrow [H\mathbb{F}_p, H\mathbb{F}_p]_{MU-mod}^*$$

and

$$l_S^* : Der_S^*(H\mathbb{F}_p, H\mathbb{F}_p) \longrightarrow [H\mathbb{F}_p, H\mathbb{F}_p]_{S-mod}^* = \mathcal{A}_p^*.$$

We already saw that the algebra of cooperations in MU -theory $\pi_* H\mathbb{F}_p \wedge_{MU} H\mathbb{F}_p$ is an exterior algebra on generators z_i . Just as the usual dual Steenrod algebra it is a Hopf algebra and it is easy to see that z_i 's are primitive elements in it. Therefore the MU -Steenrod algebra $[H\mathbb{F}_p, H\mathbb{F}_p]_{MU-mod}^*$ is itself an exterior algebra $\Lambda_{\mathbb{F}_p}(z_1^*, z_2^*, \dots)$ where z_i^* 's are dual to z_i .

Then the map l_{MU}^* sends the derivation $y_i \in Der_{MU}^{2i-1}(H\mathbb{F}_p, H\mathbb{F}_p)$ to the operation $z_i^* \in [H\mathbb{F}_p, H\mathbb{F}_p]_{MU-mod}^{2i-1}$.

Any map of MU -modules $H\mathbb{F}_p \longrightarrow \Sigma^* H\mathbb{F}_p$ can also be considered as a map of S -modules. In other words there is a map from the MU -Steenrod algebra into $\mathcal{A}_p^*$:

$$[H\mathbb{F}_p, H\mathbb{F}_p]_{MU-mod}^* \longrightarrow [H\mathbb{F}_p, H\mathbb{F}_p]_{S-mod}^* = \mathcal{A}_p^*.$$

The latter map is a map of Hopf algebras and should therefore respect primitive elements. We know that the odd degree primitive elements are the Milnor Boksteins $Q_i \in \mathcal{A}_p^*$. Since $|Q_i| = 2p^i - 1$ only the primitives $z_{2p^i-1}^*$ could have a nonzero image in $\mathcal{A}_p^*$. To see that $z_{2p^i-1}^*$ does indeed have a nonzero image notice that it is the first nonzero Postnikov invariant of the connective Morava K -theory spectrum $k(n)$. This invariant is nontrivial both in the category of S -modules and MU -modules which shows that the image of $z_{2p^i-1}^*$ is Q_i up to an invertible scalar factor.

This allows one to calculate the image of the map l_S^* . On the one hand its image should be contained in the subspace spanned by the primitive odd degree elements in $\mathcal{A}_p^*$, that is, the elements Q_i . On the other hand any operation Q_i can be lifted to a topological derivation in $Der_S^*(H\mathbb{F}_p, H\mathbb{F}_p)$ since it even lifts to an MU -derivation in $Der_{MU}^*(H\mathbb{F}_p, H\mathbb{F}_p)$. Therefore the image of l_S is the whole subspace of $\mathcal{A}_p^*$ spanned by Q_i 's.

We now turn to the calculation, due to M.Basterra and M.Mandell, of $TAQ_S^*(H\mathbb{F}_p, H\mathbb{F}_p)$. We have:

$$\begin{aligned} TAQ_S^*(H\mathbb{F}_p, H\mathbb{F}_p) &\simeq [\text{hocolim}_{n \rightarrow \infty} \Omega^n I(S^n \hat{\otimes} (H\mathbb{Z} \wedge H\mathbb{F}_p)), H\mathbb{F}_p]_{H\mathbb{F}_p-mod}^* \\ &\simeq [\text{hocolim}_{n \rightarrow \infty} \Sigma^{-n} I(S^{n-2} \hat{\otimes} [S^2 \hat{\otimes} (H\mathbb{F}_p \wedge H\mathbb{F}_p)]), H\mathbb{F}_p]_{H\mathbb{F}_p-mod}^* \\ &\simeq [\Sigma^{-2} \text{hocolim}_{n \rightarrow \infty} \Sigma^{-n+2} I(S^{n-2} \hat{\otimes} [S^2 \hat{\otimes} (H\mathbb{F}_p \wedge H\mathbb{F}_p)]), H\mathbb{F}_p]_{H\mathbb{F}_p-mod}^* \\ &\simeq \Sigma^{-2} TAQ_{H\mathbb{F}_p}^*(S^2 \hat{\otimes} (H\mathbb{F}_p \wedge H\mathbb{F}_p), H\mathbb{F}_p). \end{aligned}$$

Lemma 6.4. *There is a weak equivalence of commutative $H\mathbb{F}_p$ -algebras*

$$(6.2) \quad S^2 \hat{\otimes} (H\mathbb{F}_p \wedge H\mathbb{F}_p) \cong H\mathbb{F}_p \vee \Sigma^3 H\mathbb{F}_p.$$

Here the right hand side of (6.2) is the square-zero extension of $H\mathbb{F}_p$ by $\Sigma^3 H\mathbb{F}_p$.

To see this note first that

$$S^1 \hat{\otimes} (H\mathbb{F}_p \wedge H\mathbb{F}_p) \simeq \mathbf{THH}^S(H\mathbb{F}_p, H\mathbb{F}_p).$$

Therefore

$$\pi_* S^1 \hat{\otimes} (H\mathbb{F}_p \wedge H\mathbb{F}_p) = \mathbb{F}_p[x]$$

with $|x| = 2$, see Remark 6.2. Denote the commutative S -algebra $S^1 \hat{\otimes} (H\mathbb{F}_p \wedge H\mathbb{F}_p)$ by X . We have the following weak equivalences of commutative S -algebras:

$$S^2 \hat{\otimes} (H\mathbb{F}_p \wedge H\mathbb{F}_p) \simeq S^1 \hat{\otimes} X \simeq H\mathbb{F}_p \wedge_X H\mathbb{F}_p.$$

From the spectral sequence

$$Tor_{**}^{X*}(\mathbb{F}_p, \mathbb{F}_p) = \Lambda_{\mathbb{F}_p}(y) \Rightarrow \pi_* H\mathbb{F}_p \wedge_X H\mathbb{F}_p$$

we see that

$$\pi_* H\mathbb{F}_p \wedge_X H\mathbb{F}_p = \pi_* S^2 \hat{\otimes} (H\mathbb{F}_p \wedge H\mathbb{F}_p) = \Lambda_{\mathbb{F}_p}(y).$$

Here y has degree 3. It is not hard to see, using Proposition 2.2 that any augmented $H\mathbb{F}_p$ -algebra whose coefficient ring is an exterior algebra on one generator y in positive degree is in fact weakly equivalent to the square-zero extension $H\mathbb{F}_p \vee \Sigma^{|y|} H\mathbb{F}_p$. This finishes our sketch of the proof of Lemma 6.4.

It follows that

$$TAQ_S^*(H\mathbb{F}_p, H\mathbb{F}_p) = \Sigma^{-2} TAQ_{H\mathbb{F}_p}^*(H\mathbb{F}_p \vee \Sigma^3 H\mathbb{F}_p, H\mathbb{F}_p).$$

So we need to compute the topological Andr -Quillen cohomology of the ‘Eilenberg-MacLane object’ $H\mathbb{F}_p \vee \Sigma^3 H\mathbb{F}_p$ in the category of commutative augmented $H\mathbb{F}_p$ -algebras. There is an obvious element (‘fundamental class’) $x \in TAQ_{H\mathbb{F}_p}^3(H\mathbb{F}_p \vee \Sigma^3 H\mathbb{F}_p, H\mathbb{F}_p)$ corresponding to the identity map on $H\mathbb{F}_p \vee \Sigma^3 H\mathbb{F}_p$. Further one introduces the action of Steenrod operations on TAQ . Let $p = 2$. Then there exist operations

$$Sq^i : TAQ_{H\mathbb{F}_p}^q(-, H\mathbb{F}_p) \longrightarrow TAQ_{H\mathbb{F}_p}^{q+i}(-, H\mathbb{F}_p)$$

defined for $q < i$ and satisfying the usual Adem relations. Applying Sq^i to $y = \Sigma^{-2}x \in TAQ_{H\mathbb{F}_p}^1(H\mathbb{F}_p, H\mathbb{F}_p)$ one obtains new elements in $TAQ_{H\mathbb{F}_p}^*(H\mathbb{F}_p, H\mathbb{F}_p)$. The final result is (for $p = 2$):

Theorem 6.5. *$TAQ_S^*(H\mathbb{F}_p, H\mathbb{F}_p)$ is spanned by the elements $Sq^{s_1} Sq^{s_2} \dots Sq^{s_r} y$ where the sequence $(s_1, s_2, \dots, s_r)$ is Steenrod admissible and $s_r > 3$.*

There is a similar result for odd primes which uses the odd primary operations in TAQ instead of Steenrod squares.

To conclude this section note that the forgetful map

$$TAQ_S^*(H\mathbb{F}_p, H\mathbb{F}_p) \longrightarrow [H\mathbb{F}_p, H\mathbb{F}_p]_{S-mod}^* = \mathcal{A}_p^*$$

is not as interesting as in the associative case. It is easy to see that its image contains (multiples of) the Bokstein homomorphism β corresponding to the topological singular extension

$$H\mathbb{F}_p \longrightarrow H\mathbb{F}_{p^2} \longrightarrow H\mathbb{F}_p$$

and nothing else. Indeed, if it contained, e.g. the operation Q_i for some $i > 0$ then the corresponding singular extension $H\mathbb{F}_p \longrightarrow ? \longrightarrow \Sigma^{2(p^i-1)} H\mathbb{F}_p$ would realize the 2-stage Postnikov tower of $k(n)$ as a commutative S -algebra which is impossible.

7. EXISTENCE OF S -ALGEBRA STRUCTURES

Topological Andr -Quillen cohomology was originally introduced in [15] with the purpose of proving that BP , the Brown-Peterson spectrum supports a structure of an E_∞ -ring spectra (or, equivalently, a commutative S -algebra structure), a long-standing problem posed by P. May. Unfortunately, this problem is still open, however the problem concerning *associative* structures turned out to be much more manageable. The first result in this direction is the seminal paper of A. Robinson [30] where Morava K -theory spectra at odd primes were proved to support an A_∞ -structure. There are some difficulties in interpreting Robinson’s proof within our context, particularly the

existence of a unit seems problematic. However working in the category of MU -modules it is possible to obtain considerably stronger results. Namely, in [17] and [9] it was proved that a broad class of complex-oriented cohomology theories admit structures of S -algebras. These structures are typically non-unique and in the last section we say something about the corresponding moduli spaces. Here we describe the relevant results following [17].

Let us assume that the coefficient ring R_* of our ‘ground’ commutative S -algebra R is concentrated in even degrees. For an element $x \in R_*$ we will denote by R/x the cofibre of the map $R \xrightarrow{x} R$. Let I_* be a graded ideal generated by a (possibly infinite) regular sequence of homogeneous elements $(u_1, u_2, \dots) \in R_*$. We assume in addition that each u_k is a nonzero divisor in the ring R_* . Then we can form the R -module R/I as the infinite smash product of R/u_k . It is known by work of Strickland [32] that there is a structure of an R -ring spectrum on R/I . Clearly the coefficient ring of R/I is isomorphic to R_*/I_* where R_*/I_* is understood to be the direct limit of $R_*/(u_k, u_2, \dots, u_k)$. Let us denote the R -algebra R/I by A . Our standing assumption is that A has a structure of an R -algebra (i.e. strictly associative). This may seem a rather strong condition but, as we see shortly such a situation is quite typical. In fact P.Goerss proved in [9] that any spectrum obtained by killing a regular sequence in MU , the complex cobordism S -algebra, has a structure of an MU -algebra.

The construction we are about to describe allows one to construct new R -algebras by ‘adjoining’ the indeterminates u_k to the R -algebra A . Let us introduce the notation $A(u_k^l)_*$ for the R_* -algebra

$$\lim_{n \rightarrow \infty} R_*/(u_1, u_2, \dots, u_{k-1}, u_k^l, u_{k+1}, \dots, u_n).$$

For each l reduction modulo u_k^l determines a map of R_* -algebras

$$A(u_k^{l+1})_* \rightarrow A(u_k^l)_*.$$

Now we can formulate our main theorem in this section.

Theorem 7.1. *For each l there exist R -algebras $A(u_k^l)$ with coefficient rings $A(u_k^l)_*$ and R -algebra maps*

$$R_{l,k} : A(u_k^{l+1}) \rightarrow A(u_k^l)$$

which give the reductions mod u_k^l on the level of coefficient rings.

Let us explain the idea of the proof. One uses induction on l . The case $l = 1$ is just our original assumption that A is an R -algebra. Suppose that the R -algebras $A(u_k^l)$ with the required properties were constructed for $l \leq i$. One then shows that R/u_k^{i+1} possesses an R -ring structure and there exists a homotopy cofibre sequence of R -modules

$$(7.1) \quad R/u_k^{i+1} \longrightarrow R/u_k^i \longrightarrow \Sigma^{di+1} R/u_k$$

where the first arrow is an R -ring spectrum map realizing the reduction mod u^i in homotopy and the second arrow is an appropriate Bokstein operation. Taking the smash product of (7.1) with $R/u_1 \wedge R/u_2 \wedge \dots \wedge R/u_{k-1} \wedge R/u_{k+1} \wedge \dots$ we get the following homotopy cofibre sequence of R -modules:

$$(7.2) \quad A(u_k^{i+1}) \longrightarrow A(u_k^i) \longrightarrow \Sigma^{di+1} A.$$

Finally one proves that the sequence (7.2) could be improved to a topological singular extension by computing $Der_R^*(A(u_k^i), A)$ and analyzing the forgetful map

$$Der_R^*(A(u_k^i), A) \longrightarrow [A(u_k^i), A]_{R-mod}^*.$$

Theorem 7.1 leads to a host of examples of S -algebras. Let $R = MU_{(p)}$, the complex cobordism spectrum localized at p and $I_* = (p, x_1, x_2, \dots)$ be the maximal ideal in $MU_{(p)*} = \mathbb{Z}_{(p)}[x_1, x_2, \dots]$.

Then $MU/I \simeq H\mathbb{F}_p$ is an MU -algebra. For any polynomial generator $x_k \in MU_{(p)}$ and any i we can construct an $MU_{(p)}$ -algebra realizing in homotopy the $MU_{(p)*}$ -module $\mathbb{F}_p[x_k^i]$. Taking the homotopy inverse limit of these we get an $MU_{(p)}$ -algebra whose coefficient ring is $\mathbb{F}_p[x_k]$. Iterating this procedure we can realize topologically the $MU_{(p)*}$ -algebra $\mathbb{F}_p[x_{k_1}, x_{k_2}, \dots]$ for any sequence (possibly infinite) of polynomial generators $x_{k_1}, x_{k_2}, \dots$. We could also set $I_* = (x_1, x_2, \dots)$ to obtain integral versions of these $MU_{(p)}$ -algebras.

Recall that the Brown-Peterson spectrum BP is obtained from $MU_{(p)}$ by killing all polynomial generators in $MU_{(p)*}$ except for $v_i = x_{2(p^i-1)}$ provided that we use Hazewinkel generators for $MU_{(p)*}$. Therefore $BP_* = \mathbb{Z}_{(p)}[v_1, v_2, \dots]$. It follows that BP possesses an $MU_{(p)}$ -algebra structure. We define

$$BP\langle n \rangle = BP/(v_i, i > n) =$$

$$P(n) = BP/(v_i, i < n)$$

$$B(n) = v_n^{-1}P(n)$$

$$k(n) = BP/(v_i, i \neq n)$$

$$K(n) = v_n^{-1}BP/(v_i, i \neq n)$$

$$E(n) = v_n^{-1}BP/(v_i, i > n).$$

Note that inverting an element in the coefficient ring of an $MU_{(p)}$ -ring spectrum is an instance of Bousfield localization in the homotopy category of $MU_{(p)}$ -modules. Further Bousfield localization preserves algebra structures we conclude that all spectra listed above admit structures of $MU_{(p)}$ -algebras (and therefore MU -algebras).

Finally we need to mention the result of Hopkins and Miller, cf. [27], which states that a suitable completion E_n of the spectrum $E(n)$ (which came to be popularly known as the Morava E -theory) admits a unique structure of an S -algebra. A generalization of this theorem is discussed in [17]. Furthermore, in [10] Hopkins and Goerss proved using a more advanced technology that E_n in fact admits a unique structure of a *commutative* S -algebra.

8. SPACES OF MULTIPLICATIVE MAPS

In this section we will consider the problem of computing the homotopy type of the mapping space $\text{Map}_{C_R^{ass}}(A, B)$ where A is a cofibrant R -algebra. Unexpectedly, it turns out that the space of based loops on $\text{Map}_{C_R^{ass}}(A, B)$ is weakly equivalent to an infinite loop space. More precisely, in [18] the following theorem is proved.

Theorem 8.1. *For a q -cofibrant algebra A and a map of R -algebras $f : A \rightarrow B$ the space d -fold based loops $\Omega^d(F_{R-alg}(A, B), f)$ is weakly equivalent to the space $\Omega^\infty \mathbf{Der}_R(A, \Sigma^{-d}B)$.*

The main ingredient in the proof of this theorem is the following

Lemma 8.2. *There is a weak equivalence of R -algebras A^{S^d} and the square-zero-extension $A \vee \Sigma^{-d}A$.*

Note that, obviously, there is a weak equivalence of R -modules

$$(8.1) \quad A^{S^d} \longrightarrow A \vee \Sigma^{-d}A$$

The lemma is proved by computing the topological derivations of A^{S^d} and showing that the map of (8.1) could be improved to a topological derivation of A^{S^d} with values in $\Sigma^{-d}A$.

The proof of Theorem 8.1 is actually very easy. We have the following commutative diagram of spaces where both rows are homotopy fibre sequences:

$$\begin{array}{ccccc}
 \mathcal{Top}_*(S^d, \text{Map}_{\mathcal{C}_R^{ass}}(A, B)) & \longrightarrow & \mathcal{Top}(S^d, \text{Map}_{\mathcal{C}_R^{ass}}(A, B)) & \longrightarrow & \text{Map}_{\mathcal{C}_R^{ass}}(A, B) \\
 \downarrow & & \downarrow & & \parallel \\
 ? & \longrightarrow & \text{Map}_{\mathcal{C}_R^{ass}}(A, B^{S^d}) & \longrightarrow & \text{Map}_{\mathcal{C}_R^{ass}}(A, B)
 \end{array}$$

Here the horizontal rightmost arrows are both induced by the inclusion of the base point into S^d . Since the right and the middle vertical arrows are weak equivalences (even isomorphisms) it follows that the map $\mathcal{Top}_*(S^d, \text{Map}_{\mathcal{C}_R^{ass}}(A, B)) \rightarrow ?$ is a weak equivalence. But Lemma 8.2 tells us that the R -algebra B^{S^d} is weakly equivalent as an R -algebra to $B \vee \Sigma^{-d}B$. In other words the term $?$ is weakly equivalent to the topological space of maps $A \rightarrow B \vee \Sigma^{-d}B$ which yield f when composed with the projection $B \vee \Sigma^{-d}B \rightarrow B$. Therefore $?$ is weakly equivalent to $\Omega^\infty \mathbf{Der}_R(A, \Sigma^{-d}B)$ and our theorem is proved.

Corollary 8.3. *For a q -cofibrant algebra A and a map of R -algebras $f : A \rightarrow B$ there is a bijection between sets $\pi_d(\text{Map}_{\mathcal{C}_R^{ass}}(A, B), f)$ and $\text{Der}_R^{-d}(A, B)$ for $d \geq 1$. If $d \geq 2$ then this bijection is an isomorphism of abelian groups.*

One might wonder whether Theorem 8.1 remains true in the context of commutative S -algebras. The answer is no. The crucial point is the weak equivalence of S -algebras $S \vee S^{-1}$ and S^{S^1} . It is clear that $\pi_0 S \wedge_{S \vee S^{-1}} S$ is the divided power ring. However N.Kuhn and M.Mandell proved that $\pi_0 S \wedge_{S^{S^1}} S$ is the ring of numeric polynomials. Therefore $S \vee S^{-1}$ and S^{S^1} cannot be weakly equivalent as commutative S -algebras.

Another point of view is afforded by Mandell's theorem [21] which states that the homotopy category of connected p -complete nilpotent spaces of finite p -type can be embedded into the category of E_∞ algebras over $\mathbb{F}_p$ as a full subcategory. Here $\mathbb{F}_p$ is the algebraic closure of the field $\mathbb{F}_p$. This embedding is via the singular cochains functor $C^*(-, \mathbb{F}_p)$. The category of E_∞ algebras over $\mathbb{F}_p$ is Quillen equivalent to the category of commutative $H\mathbb{F}_p$ -algebras. Consider an arbitrary space X (connected, p -complete, nilpotent and of finite p -type) and the corresponding $H\mathbb{F}_p$ -algebra $\tilde{X}$ which we assume to be cofibrant. Then the space of maps in $\mathcal{C}_{H\mathbb{F}_p}^{comm}$ from $\tilde{X}$ into $H\mathbb{F}_p$ will be weakly equivalent to X . This shows that the space of commutative R -algebra maps could have an essentially arbitrary homotopy type.

However, there is a context in which the analogue of Theorem 8.1 holds for commutative R -algebras. Let us suppose that our ground S -algebra R is rational, that is, local with respect to $H\mathbb{Q}$. Let A and B be commutative R -algebras where A is cofibrant as before. Then it is easy to see that A^{S^1} and $A \vee \Sigma^{-1}A$ are weakly equivalent as *commutative* R -algebras. Indeed, $A \vee \Sigma^{-1}A$ is weakly equivalent as a commutative R -algebra $A \wedge R[S^{-1}]$ where $R[S^{-1}]$ is the free commutative R -algebra on the R -module S^{-1} . We have $\pi_* A^{S^1} = \Lambda_{A_*}(x)$ where the exterior generator x has degree -1 . The element x determines a map of R -modules $S^1 \rightarrow A^{S^1}$ and therefore a map of R -algebras $A \vee \Sigma^{-1}A \simeq A \wedge R[S^{-1}] \rightarrow A^{S^1}$ which is a weak equivalence. Further there are the following equivalences of commutative R -algebras:

$$A^{S^d} \cong A^{S^1 \wedge S^{d-1}} \cong (A^{S^1})^{S^{d-1}} \simeq (A \vee \Sigma^{-1}A)^{S^{d-1}} \simeq A \vee \Sigma^{-d}A.$$

So we in this case we have the exact analogue of Lemma 8.2. The remainder of the proof is the same and we obtain

Theorem 8.4. *Let R be a rational commutative S -algebra, Then for a q -cofibrant R -algebra A and a map of R -algebras $f : A \rightarrow B$ the space d -fold based loops $\Omega^d(\text{Map}_{\mathcal{C}_R^{comm}}, f)$ is weakly equivalent to*

the space $\Omega^\infty \mathbf{TAQ}_R(A, \Sigma^{-d}B)$. Furthermore there is a bijection between sets $\pi_d(\mathrm{Map}_{\mathcal{C}_R^{\mathrm{comm}}}(A, B), f)$ and $TAQ_R^{-d}(A, B)$ for $d \geq 1$. If $d \geq 2$ then this bijection is an isomorphism of abelian groups.

For $R = H\mathbb{Q}$, the Eilenberg-MacLane spectrum the category of commutative R -algebras is Quillen equivalent to the category of commutative differential graded algebras over $\mathbb{Q}$. Via this equivalence Theorem 8.4 translates into a statement in rational homotopy theory which does not seem to have appeared in the literature.

9. MODULI SPACES

For an R -module X it is natural to look at the set of all nonisomorphic R -algebra structures on X . (We assume that X supports at least one R -algebra structure.) This set is actually π_0 of the Dwyer-Kan *classification space*, cf. [6] which we will now describe.

Consider the subcategory W of $\mathcal{C}_R^{\mathrm{ass}}$ consisting of those R -algebras which are weakly equivalent to X as R -modules. The morphisms of W are the weak equivalences of R -algebras. Then the *moduli space* $\mathcal{M}(X)$ is by definition the nerve of W .

For a cofibrant R -algebra A we will denote by $haut(A)$ the topological monoid of homotopy autoequivalences of A . For a noncofibrant A we take for $haut(A)$ the monoid of homotopy autoequivalences of its cofibrant approximation. Thus $haut(A)$ is well defined up to homotopy. We can form its classifying space $Bhaut(A)$. Then the results of Dwyer and Kan imply the following

Theorem 9.1. *There is a weak equivalence of spaces*

$$\mathcal{M}(X) \simeq \coprod B\mathrm{haut}(A)$$

where the disjoint union is taken over the set of connected components of W .

We assumed from the beginning the existence of at least one R -algebra structure on X , that gives the space $\mathcal{M}(X)$ a base point. Let us denote the corresponding R -algebra by A . The connected component of A in $\mathcal{M}(X)$ is, therefore, weakly equivalent to $B\mathrm{haut}(A)$.

Note that it is possible to consider other moduli spaces for X . For example, we could insist that the homotopy X_* has a fixed ring structure, or that X itself has a fixed R -ring spectrum structure up to homotopy. Each of these choices affects π_0 of our moduli space but not the connected components themselves, and therefore, not the higher homotopy groups.

Assume, without loss of generality, that A is a cofibrant R -algebra. Clearly $haut(A)$ is a disjoint union summand of the space $\mathrm{Map}_{\mathcal{C}_R^{\mathrm{ass}}}(A, A)$ of all multiplicative self-maps of A . Taking the identity map $A \rightarrow A$ as the base point in both $haut(A)$ and $\mathrm{Map}_{\mathcal{C}_R^{\mathrm{ass}}}(A, A)$ we see that

$$\Omega \mathrm{haut}(A) \simeq \Omega \mathrm{Map}_{\mathcal{C}_R^{\mathrm{ass}}}(A, A).$$

But we know from the previous section that there is a weak equivalence of spaces

$$\Omega \mathrm{Map}_{\mathcal{C}_R^{\mathrm{ass}}}(A, A) \simeq \Omega \Omega^\infty \mathbf{Der}_R(A, A).$$

This gives the following

Theorem 9.2. *The space $\Omega^2 \mathcal{M}(X)$ of two-fold loops on the moduli space $\mathcal{M}(X)$ is weakly equivalent to $\Omega \Omega^\infty \mathbf{Der}_R(A, A)$. In particular it is an infinite loop space.*

In particular, we have an isomorphism

$$\pi_i \mathcal{M}(X) \cong \mathrm{Der}_R^{1-i}(A, A)$$

for $i \geq 2$. The computation of π_1 and π_0 is, of course, a completely different story.

What about the moduli space of *commutative* R -algebra structures? Theorem 9.1 has an obvious analogue in the context of commutative R -algebras. However the analogue of Theorem 9.2 falls through because for a commutative R -algebra A the space of self-maps of A in $\mathcal{C}_R^{\mathrm{comm}}$ is not in

any obvious way related to $\mathbf{TAQ}_R(A, A)$. However Hopkins and Goerss proved in [10] that E_n , the Morava E -theory spectrum mentioned at the end of Section 7 admits a unique structure of a commutative S -algebra and the space of its self-maps is homotopically discrete with the set of connected components being equal to S_n , the Morava stabilizer group. In our interpretation it means that the appropriate moduli space of E_n is weakly equivalent to BS_n , the classifying space of S_n .

REFERENCES

- [1] M. Bastera, *André-Quillen cohomology of commutative S -algebras*. J. Pure Appl. Algebra 144 (1999), no. 2, 111–143.
- [2] M. Bastera & R. McCarthy, *Γ -homology, topological André-Quillen homology and stabilization*. Topology Appl. 121 (2002), no.3, 551–566.
- [3] A. Baker & A. Lazarev, *Topological Hochschild cohomology and generalized Morita equivalence*, preprint, math.AT/0209003.
- [4] M. Bökstedt, *Topological Hochschild homology of $\mathbb{Z}$ and $\mathbb{Z}_p$* , preprint.
- [5] J. Cuntz & D. Quillen, *Algebra extensions and nonsingularity*. J. Amer. Math. Soc. 8 (1995), no. 2, 251–289.
- [6] W. Dwyer & D. Kan, *A classification theorem for diagrams of simplicial sets*. Topology 23 (1984), no. 2, 139–155.
- [7] A. Elmendorf, I. Kriz, M. Mandell & J. P. May, *Rings, modules, and algebras in stable homotopy theory*. Mathematical Surveys and Monographs **47** (1997).
- [8] V. Franjou & J. Lannes, L. Schwartz, *Autour de la cohomologie de Mac Lane des corps finis*. Invent. Math. 115 (1994), no. 3, 513–538.
- [9] P. Goerss, *Associative MU -algebras*, preprint 2001.
- [10] P. Goerss & M. Hopkins *Realizing commutative ring spectra as E_∞ ring spectra*, preprint.
- [11] T. Goodwillie, *Calculus I. The first derivative of pseudoisotopy theory*. K-Theory 4 (1990), no. 1, 1–27.
- [12] T. Goodwillie, *Calculus 3. The Taylor series of homotopy functors*, preprint.
- [13] M. Hovey, *Model categories*. Mathematical Surveys and Monographs, 63. American Mathematical Society, Providence, RI, 1999.
- [14] M. Hovey, B. Shipley & J. Smith, *Symmetric spectra*. J. Amer. Math. Soc. 13 (2000), no. 1, 149–208.
- [15] I. Kriz, *Towers of E_∞ -ring spectra with applications to BP* , preprint, 1993.
- [16] A. Lazarev, *Homotopy theory of A_∞ ring spectra and applications to MU -modules*. K-Theory 24 (2001), no. 3, 243–281.
- [17] A. Lazarev, *Towers of MU -algebras and a generalized Hopkins-Miller theorem*, to appear in the Proc. of the London Math. Soc.
- [18] A. Lazarev, *Spaces of multiplicative maps between highly structured ring spectra*, Proceedings of the Isle of Sky Conference on Algebraic Topology, to appear.
- [19] A. Lindenstrauss & I. Madsen, *Topological Hochschild homology of number rings*. Trans. Amer. Math. Soc. 352 (2000), no. 5, 2179–2204.
- [20] I. Madsen, *Algebraic K -theory and traces*. Current developments in mathematics, 1995 (Cambridge, MA), 191–321, Internat. Press, Cambridge, MA, 1994.
- [21] M. Mandell, *E_∞ algebras and p -adic homotopy theory*. Topology 40 (2001), no. 1, 43–94.
- [22] J. P. May, *E_∞ ring spaces and E_∞ ring spectra*. With contributions by F. Quinn, N. Ray, and J. Tornehave. Lecture Notes in Mathematics, Vol. 577. Springer-Verlag, Berlin-New York, 1977.
- [23] J. P. May, *The geometry of iterated loop spaces*. Lectures Notes in Mathematics, Vol. 271. Springer-Verlag, Berlin-New York, 1972.
- [24] J. McClure & R. Staffeldt, *On the topological Hochschild homology of bu* . I. Amer. J. Math. 115 (1993), no. 1, 1–45.
- [25] T. Pirashvili & B. Richter, *Robinson-Whitehouse complex and stable homotopy*. Topology 39 (2000), no. 3, 525–530.
- [26] D. Quillen, *Homotopical algebra*. Lecture Notes in Mathematics, No. 43. Springer-Verlag, Berlin-New York 1967.
- [27] C. Rezk, *Notes on the Hopkins-Miller Theorem. Homotopy theory via algebraic geometry and group representations* (Evanston, IL, 1997), 313–366, Contemp. Math., 220, Amer. Math. Soc., Providence, RI, 1998.
- [28] B. Richter, *An Atiyah-Hirzebruch spectral sequence for topological André-Quillen homology*. J. Pure Appl. Algebra 171 (2002), no. 1, 59–66.
- [29] A. Robinson & S. Whitehouse, *Operads and Γ -homology of commutative rings*. Math. Proc. Cambridge Philos. Soc. 132 (2002), no. 2, 197–234.
- [30] A. Robinson, *Obstruction theory and the strict associativity of Morava K -theories*. Advances in homotopy theory 143–152, London Math. Soc. Lecture Note Ser., 139, Cambridge Univ. Press, Cambridge, 1989.

- [31] A.Robinson, *Gamma homology, Lie representations and E_∞ multiplications*, preprint available at <http://www.maths.warwick.ac.uk/~car/>
- [32] N.P.Strickland, *Products on MU-modules*. Trans. Amer. Math. Soc. 351 (1999), no. 7, 2569–2606.

MATHEMATICS DEPARTMENT, BRISTOL UNIVERSITY, BRISTOL, BS8 1TW, ENGLAND.
E-mail address: `a.lazarev@bristol.ac.uk`